

ADDITION FORMULA FOR 2-PARAMETER FAMILY OF ASKEY-WILSON POLYNOMIALS

H.T. KOELINK

December 1994

ABSTRACT. For a two parameter family of Askey-Wilson polynomials, that can be regarded as basic analogues of the Legendre polynomials, an addition formula is derived. The addition formula is a two-parameter extension of Koornwinder's addition formula for the little q -Legendre polynomials. A corresponding product formula is derived. As the base tends to one, the addition and product formula go over into the addition and product formula for the Legendre polynomial. The addition formula is derived from the interpretation of Askey-Wilson polynomials as generalised matrix elements on the quantum $SU(2)$ group.

1. INTRODUCTION

Let us start with describing the classical addition formula for the Legendre polynomials. The Jacobi polynomials $R_n^{(\alpha, \beta)}(x)$ are orthogonal polynomials with respect to the beta measure $(1-x)^\alpha(1+x)^\beta$ on the interval $[-1, 1]$ and normalised by $R_n^{(\alpha, \beta)}(1) = 1$. The polynomial $R_n^{(0,0)}(x)$ is the Legendre polynomial, and for $\alpha = \beta$ the name Gegenbauer or ultraspherical polynomials is used. Laplace proved the following addition formula for the Legendre polynomials in 1782;

$$(1.1) \quad \begin{aligned} & R_l^{(0,0)}(xy + t\sqrt{(1-x^2)(1-y^2)}) = R_l^{(0,0)}(x)R_l^{(0,0)}(y) \\ & + 2 \sum_{n=1}^l \frac{(l+n)!}{(l-n)!(n!)^2} 2^{-2n} (1-x^2)^{n/2} R_{l-n}^{(n,n)}(x) (1-y^2)^{n/2} R_{l-n}^{(n,n)}(y) T_n(t), \end{aligned}$$

where $T_n(\cos \theta) = \cos n\theta = R_n^{(-1/2, -1/2)}(\cos \theta)$, is the Chebyshev polynomial of the first kind, cf. Askey [2, Lecture 4]. The orthogonality relations for the Chebyshev polynomials

1991 *Mathematics Subject Classification.* 33D45, 33D80.

Key words and phrases. Askey-Wilson polynomials, big q -Jacobi polynomials, addition formula, product formula, quantum group, $SU(2)$.

Supported by a Fellowship of the Research Council of the Katholieke Universiteit Leuven.

yield the product formula

$$(1.2) \quad R_{l-n}^{(n,n)}(x)R_{l-n}^{(n,n)}(y) = 2^{2n} \frac{(l-n)!(n!)^2}{\pi(l+n)!} (\sqrt{(1-x^2)(1-y^2)})^{-n} \\ \times \int_{-1}^1 R_l^{(0,0)}(xy + t\sqrt{(1-x^2)(1-y^2)}) \frac{T_n(t)}{\sqrt{1-t^2}} dt,$$

which for $n = 0$ is the product formula for the Legendre polynomial.

For basic analogues of the Legendre polynomials there are several analogues of the addition formula (1.1). The addition formula for the continuous q -Legendre polynomials is proved by Rahman and Verma [23] as a special case of their addition formula for the continuous q -ultraspherical polynomials, which have been introduced by Rogers in 1895. A quantum $SU(2)$ group theoretic proof of the addition formula for the continuous q -Legendre polynomials has been given by the author [9]. Also, the quantum $SU(2)$ group has naturally led to an addition formula for the little q -Legendre polynomials, cf. Koornwinder [15], since the matrix elements of the irreducible unitary representations of the quantum $SU(2)$ group are explicitly known in terms of little q -Jacobi polynomials [24], [17], [13]. See Rahman [22] for an analytic proof of this addition formula. There is also an addition formula for the big q -Legendre polynomials [10], and the addition formula derived in this paper contains these last two addition formulas as limiting cases. Apart from the quantum group theoretic approach it is also possible to use quantum algebras and its representation theory in proving addition formulas for q -special functions, see e.g. Kalnins and Miller [8] and references therein.

As a result of Koornwinder's work [16] on zonal spherical elements on the quantum $SU(2)$ group and Askey-Wilson polynomials, there have been papers by Noumi and Mimachi [19], [20] and the author [9], [11], in which the full four parameter family of Askey-Wilson polynomials is interpreted as generalised matrix elements on the quantum $SU(2)$ group. This has led to an abstract addition formula for Askey-Wilson polynomials, i.e. involving non-commuting variables, [19], [20], [9], [11], from which a (degenerate) addition formula for Askey-Wilson polynomials can be obtained.

In this paper we derive an addition formula for a two-parameter family of Askey-Wilson polynomials much along the lines of the quantum group theoretic proof of the addition formula for the little q -Legendre polynomials. The proof is based on the fact that we have a basis of eigenvectors for a certain self-adjoint operator in an irreducible $*$ -representation of the C^* -algebra for the quantum $SU(2)$ group [10] and that the non-polynomial minimal part of the associated spherical generalised elements can be factorised in elements which act nicely in this basis of eigenvectors. This last part is suggested by the paper by Noumi and Mimachi [21], and we make an explicit identification with their work in remark 3.5. All this is explained in §3. The proof of the addition formula is then relatively easy and is given in §4. In §5 the corresponding product formula is derived and the limit transition $q \uparrow 1$ to (1.1) and (1.2) is considered in §6. In §2 we recall the necessary results on the relation between certain basic hypergeometric orthogonal polynomials and the quantum $SU(2)$ group.

To end this introduction we remark that this paper is concerned with transforming an identity for q -special functions involving non-commuting variables into an identity for

q -special functions in commuting variables.

2. ORTHOGONAL POLYNOMIALS AND THE QUANTUM $SU(2)$ GROUP

In this section we present some of the relations between basic hypergeometric orthogonal polynomials and the quantum $SU(2)$ group that are necessary to obtain an addition formula for a two-parameter family of Askey-Wilson polynomials. More information can be found in the survey papers by Koornwinder [14], Noumi [18] and the author [11].

$\mathcal{A}_q(SU(2))$ is the complex unital associative algebra generated by $\alpha, \beta, \gamma, \delta$ subject to the relations

$$(2.1) \quad \begin{aligned} \alpha\beta &= q\beta\alpha, & \alpha\gamma &= q\gamma\alpha, & \beta\delta &= q\delta\beta, & \gamma\delta &= q\delta\gamma, \\ \beta\gamma &= \gamma\beta, & \alpha\delta - q\beta\gamma &= \delta\alpha - q^{-1}\beta\gamma = 1 \end{aligned}$$

for some constant $q \in \mathbb{C}$. With a $*$ -operator given by

$$(2.2) \quad \alpha^* = \delta, \quad \beta^* = -q\gamma, \quad \gamma^* = -q^{-1}\beta, \quad \delta^* = \alpha$$

the algebra $\mathcal{A}_q(SU(2))$ becomes a $*$ -algebra for non-zero real q , and from now on we fix $0 < q < 1$. The $*$ -algebra can be completed into a C^* -algebra, and then this quantum group has been introduced by Woronowicz [26] as an example of his general theory of compact matrix quantum groups [27].

All $*$ -representations of the C^* -algebra have been classified, cf. Vaksman and Soibelman [24], and we only use the infinite dimensional $*$ -representation π of $\mathcal{A}_q(SU(2))$ in the Hilbert space $\ell^2(\mathbb{Z}_+)$ with orthonormal basis $\{e_n \mid n \in \mathbb{Z}_+\}$ given by

$$(2.3) \quad \begin{aligned} \pi(\alpha)e_n &= \sqrt{1 - q^{2n}}e_{n-1}, & \pi(\beta)e_n &= -q^{n+1}e_n, \\ \pi(\gamma)e_n &= q^n e_n, & \pi(\delta)e_n &= \sqrt{1 - q^{2n+2}}e_{n+1}, \end{aligned}$$

where we use the convention $e_{-p} = 0$ for $p \in \mathbb{N}$. Note that $-q\pi(\gamma) = \pi(\beta)$. For all $\xi \in \mathcal{A}_q(SU(2))$, $\pi(\xi)$ is a bounded operator on $\ell^2(\mathbb{Z}_+)$.

The algebra $\mathcal{A}_q(SU(2))$ is an example of a Hopf $*$ -algebra. The comultiplication Δ , which is a $*$ -homomorphism of $\mathcal{A}_q(SU(2)) \rightarrow \mathcal{A}_q(SU(2)) \otimes \mathcal{A}_q(SU(2))$, is given on the generators by

$$(2.4) \quad \begin{aligned} \Delta(\alpha) &= \alpha \otimes \alpha + \beta \otimes \gamma, & \Delta(\beta) &= \alpha \otimes \beta + \beta \otimes \delta, \\ \Delta(\gamma) &= \gamma \otimes \alpha + \delta \otimes \gamma, & \Delta(\delta) &= \gamma \otimes \beta + \delta \otimes \delta, \end{aligned}$$

The representations of the quantum $SU(2)$ group correspond to the corepresentations of the Hopf $*$ -algebra $\mathcal{A}_q(SU(2))$. The irreducible unitary representations of the quantum $SU(2)$ group are completely classified and, as for the Lie group $SU(2)$, they are indexed by the spin l , $l \in \frac{1}{2}\mathbb{Z}_+$, for each dimension $2l+1$, cf. Woronowicz [26], [27], Koornwinder [13], Masuda et al. [17], Vaksman and Soibelman [24]. The matrix elements $t_{n,m}^l \in \mathcal{A}_q(SU(2))$, $n, m = -l, -l+1, \dots, l$, of the irreducible unitary corepresentation t^l of spin l are explicitly

known in terms of the little q -Jacobi polynomials [13], [17], [24]. The cohomorphism property for the matrix elements is

$$(2.5) \quad \Delta(t_{n,m}^l) = \sum_{k=-l}^l t_{n,k}^l \otimes t_{k,m}^l,$$

and for $n = m = 0$ Koornwinder [15] has transformed this identity into an addition formula for the little q -Legendre polynomials.

It is possible to extend the notion of matrix elements to get generalised matrix elements, which satisfy relative left and right infinitesimal invariance. In the spherical case this has been introduced by Koornwinder [14], [16] and for the associated spherical elements it has been worked out by the author [9] and for generalised matrix elements Noumi and Mimachi [19], [20] have stated the corresponding theorems, see the survey [11] for a complete proof. To describe the infinitesimal invariance the duality of the Hopf $*$ -algebra with the quantised universal enveloping algebra $\mathcal{U}_q(\mathfrak{su}(2))$ as introduced by Jimbo [7] is needed, but we only need some of the results, see [11] for proofs. There exist elements $b_{i,j}^l(\tau, \sigma) \in \mathcal{A}_q(SU(2))$, $l \in \frac{1}{2}\mathbb{Z}_+$, $i, j \in \{-l, -l+1, \dots, l-1, l\}$, $\sigma, \tau \in \mathbb{R} \cup \{\infty\}$ such that

$$(2.6) \quad \Delta(b_{i,j}^l(\tau, \sigma)) = \sum_{n=-l}^l (D.b_{i,n}^l(\tau, \mu)) \otimes b_{n,j}^l(\mu, \sigma), \quad \forall \mu \in \mathbb{R} \cup \{\infty\}.$$

Here $D.: \mathcal{A}_q(SU(2)) \rightarrow \mathcal{A}_q(SU(2))$ is the bijective algebra homomorphism given by $D.\alpha = q^{-\frac{1}{2}}\alpha$, $D.\beta = q^{\frac{1}{2}}\beta$, $D.\gamma = q^{-\frac{1}{2}}\gamma$ and $D.\delta = q^{\frac{1}{2}}\delta$. For $\sigma = \tau = \infty$ the generalised matrix element $b_{i,j}^l(\tau, \sigma)$ reduces to $t_{i,j}^l$ up to a constant, and $\sigma = \tau = \mu = \infty$ in (2.6) reduces to (2.5). The case of most interest to us is $l \in \mathbb{Z}_+$, $i = j = 0$ and $\mu = \infty$ in (2.6); i.e.

$$(2.7) \quad \Delta(b_{0,0}^l(\tau, \sigma)) = \sum_{n=-l}^l (D.b_{0,n}^l(\tau, \infty)) \otimes b_{n,0}^l(\infty, \sigma).$$

The generalised matrix elements are explicitly known in terms of Askey-Wilson polynomials, where the argument is a simple element of the Hopf $*$ -algebra $\mathcal{A}_q(SU(2))$, cf. [14], [16], [19], [20], [9], [11]. Define in $\mathcal{A}_q(SU(2))$

$$(2.8) \quad \begin{aligned} \rho_{\tau,\sigma} = \frac{1}{2} & (\alpha^2 + \delta^2 + q\gamma^2 + q^{-1}\beta^2 + i(q^{-\sigma} - q^{\sigma})(q\delta\gamma + \beta\alpha) \\ & - i(q^{-\tau} - q^{\tau})(\delta\beta + q\gamma\alpha) + (q^{-\sigma} - q^{\sigma})(q^{-\tau} - q^{\tau})\beta\gamma), \end{aligned}$$

then

$$(2.9) \quad b_{0,0}^l(\tau, \sigma) = q^{-l}(q^{2l+2}; q^2)_l^{-1} p_l^{(0,0)}(\rho_{\tau,\sigma}; q^{\tau}, q^{\sigma} \mid q^2).$$

The normalisation has been chosen differently from [11]. It is chosen such that, with (2.12), the common constant in (2.7) is dropped. Here we use the notation

$$(2.10) \quad p_n^{(\alpha,\beta)}(x; s, t \mid q) = p_n(x; q^{1/2+\alpha}s/t, q^{1/2}t/s, -q^{1/2}/(st), -stq^{1/2+\beta} \mid q)$$

for q -analogues of the Jacobi polynomial, where

$$(2.11) \quad p_n(\cos \theta; a, b, c, d \mid q) = a^{-n} (ab, ac, ad; q)_n {}_4\varphi_3 \left(\begin{matrix} q^{-n}, abcdq^{n-1}, ae^{i\theta}, a^{-i\theta} \\ ab, ac, ad \end{matrix}; q, q \right)$$

denotes an Askey-Wilson polynomial [3]. The notation for basic hypergeometric series follows the excellent book [6] by Gasper and Rahman;

$${}_{r+1}\varphi_r \left(\begin{matrix} a_1, \dots, a_{r+1} \\ b_1, \dots, b_r \end{matrix}; q, z \right) = \sum_{k=0}^{\infty} \frac{(a_1, \dots, a_{r+1}; q)_k}{(q, b_1, \dots, b_r; q)_k} z^k,$$

where the q -shifted factorials are defined by

$$(a_1, \dots, a_r; q)_k = \prod_{i=1}^r (a_i; q)_k, \quad (a; q)_k = \prod_{i=0}^{k-1} (1 - aq^i).$$

The empty product equals 1 by definition. Since $0 < q < 1$, $(a; q)_{\infty}$ is well-defined.

The other generalised matrix elements are also explicitly known in terms of the Askey-Wilson polynomials, cf. [19], [20], [9], [11], and we need the following special cases;

$$(2.12) \quad \begin{aligned} b_{n,0}^l(\infty, \sigma) &= C_n(\sigma) (\delta_{\infty, \sigma-1} \gamma_{\infty, \sigma})^n P_{l-n}^{(n,n)}(\rho_{\infty, \sigma}; q^{2\sigma}, 1; q^2), \\ b_{0,n}^l(\tau, \infty) &= C_n(\tau) (\delta_{\tau-1, \infty} \beta_{\tau, \infty})^n P_{l-n}^{(n,n)}(\rho_{\tau, \infty}; q^{2\tau}, 1; q^2), \\ b_{-n,0}^l(\infty, \sigma) &= C_n(\sigma) (\beta_{\infty, \sigma-1} \alpha_{\infty, \sigma})^n P_{l-n}^{(n,n)}(\rho_{\infty, \sigma}; q^{2\sigma+2n}, q^{2n}; q^2), \\ b_{0,-n}^l(\tau, \infty) &= C_n(\tau) (\gamma_{\tau-1, \infty} \alpha_{\tau, \infty})^n P_{l-n}^{(n,n)}(\rho_{\tau, \infty}; q^{2\tau+2n}, q^{2n}; q^2) \end{aligned}$$

with the constant given by

$$C_n(\sigma) = q^{\frac{1}{2}(l-n)(l-n-1)} q^{-\sigma l} \frac{(q^{2n+2}, -q^{2\sigma-2l}; q^2)_{l-n}}{\sqrt{(q^2, q^{2l+2n+2}; q^2)_{l-n}}}.$$

To explain the notation in (2.12) we have the following limit cases of $\rho_{\tau, \sigma}$;

$$(2.13) \quad \begin{aligned} \rho_{\tau, \infty} &= \lim_{\sigma \rightarrow \infty} 2q^{\sigma+\tau-1} \rho_{\tau, \sigma} = iq^{\tau-1} (q\delta\gamma + \beta\alpha) + q^{-1} (1 - q^{2\tau}) \beta\gamma, \\ \rho_{\infty, \sigma} &= \lim_{\tau \rightarrow \infty} 2q^{\sigma+\tau-1} \rho_{\tau, \sigma} = -iq^{\sigma-1} (\delta\beta + q\gamma\alpha) + q^{-1} (1 - q^{2\sigma}) \beta\gamma. \end{aligned}$$

The elements $\alpha_{\tau, \infty}$, $\alpha_{\infty, \sigma}$, etcetera, are the straightforward limit cases of the elements defined by

$$(2.14) \quad \begin{aligned} \alpha_{\tau, \sigma} &= q^{1/2} \alpha - iq^{\sigma-1/2} \beta + iq^{\tau+1/2} \gamma + q^{\sigma+\tau-1/2} \delta, \\ \beta_{\tau, \sigma} &= -q^{\sigma+1/2} \alpha - iq^{-1/2} \beta - iq^{\sigma+\tau+1/2} \gamma + q^{\tau-1/2} \delta, \\ \gamma_{\tau, \sigma} &= -q^{\tau+1/2} \alpha + iq^{\tau+\sigma-1/2} \beta + iq^{1/2} \gamma + q^{\sigma-1/2} \delta, \\ \delta_{\tau, \sigma} &= q^{\tau+\sigma+1/2} \alpha + iq^{\tau-1/2} \beta - iq^{\sigma+1/2} \gamma + q^{-1/2} \delta. \end{aligned}$$

These elements are up to a constant factor the generalised matrix elements $b_{i,j}^{1/2}(\tau, \sigma)$, cf. [11, prop. 6.5]. The $*$ -operator on these simple generalised matrix elements is given by (2.15)

$$\alpha_{\tau,\sigma}^* = q\delta_{\tau-1,\sigma-1}, \quad \beta_{\tau,\sigma}^* = -\gamma_{\tau-1,\sigma+1}, \quad \gamma_{\tau,\sigma}^* = -\beta_{\tau+1,\sigma-1}, \quad \delta_{\tau,\sigma}^* = q^{-1}\alpha_{\tau+1,\sigma+1}.$$

Finally, the polynomials involved in (2.12) are the big q -Jacobi polynomials introduced by Andrews and Askey [1];

$$(2.16) \quad P_n^{(\alpha,\beta)}(x; c, d; q) = {}_3\phi_2 \left(\begin{matrix} q^{-n}, q^{n+\alpha+\beta+1}, q^{1+\alpha}x/c \\ q^{\alpha+1}, -q^{1+\alpha}d/c \end{matrix}; q, q \right).$$

The special case (2.12) is obtained from the general expression for the generalised matrix elements $b_{i,j}^l(\tau, \sigma)$ in terms of Askey-Wilson polynomials by use of the limit transition of the Askey-Wilson polynomials to the big q -Jacobi polynomials as described by Koornwinder [16, (6.2)]. Equation (2.12) has been obtained by Noumi and Mimachi [21, thm. 3.5] in the context of a slightly different algebra and we make a precise identification with the algebra considered in [21] in remark 3.5.

3. BASIS OF THE REPRESENTATION SPACE

The standard orthonormal basis $\{e_n \mid n \in \mathbb{Z}_+\}$ of the representation space $\ell^2(\mathbb{Z}_+)$ of the irreducible $*$ -representation of $\mathcal{A}_q(SU(2))$ is not well suited to calculate the action of the operators $\pi(b_{n,0}^l(\infty, \sigma))$. In this section we show that these operators act nicely in a suitable basis of $\ell^2(\mathbb{Z}_+)$, which has already been introduced in [10].

Proposition 3.1. ([10, prop. 4.1, cor. 4.2]) *$\ell^2(\mathbb{Z}_+)$ has an orthogonal basis of eigenvectors $v_\lambda(q^\sigma)$, where $\lambda = -q^{2n}$, $n \in \mathbb{Z}_+$, and $\lambda = q^{2\sigma+2n}$, $n \in \mathbb{Z}_+$, for the eigenvalue λ of the self-adjoint operator $\pi(\rho_{\infty,\sigma})$. Moreover,*

$$\begin{aligned} \langle v_\lambda(q^\sigma), v_\lambda(q^\sigma) \rangle &= q^{-2n}(q^2; q^2)_n(-q^{2-2\sigma}; q^2)_n(-q^{2\sigma}; q^2)_\infty, & \lambda = -q^{2n}, \\ \langle v_\lambda(q^\sigma), v_\lambda(q^\sigma) \rangle &= q^{-2n}(q^2; q^2)_n(-q^{2+2\sigma}; q^2)_n(-q^{-2\sigma}; q^2)_\infty, & \lambda = q^{2\sigma+2n}. \end{aligned}$$

Remark 3.2. Koornwinder [16], see also Noumi and Mimachi [21], has proved that the Haar functional on the commutative $*$ -subalgebra of $\mathcal{A}_q(SU(2))$ generated by $\rho_{\infty,\sigma}$ is given by $h(p(\rho_{\infty,\sigma})) = \int_{-1}^{q^{2\sigma}} p(x) d_{q^2}x$, see [6, §1.11] for the definition of a q -integral. It is remarkable that the spectrum of the self-adjoint operator $\pi(\rho_{\infty,\sigma})$ equals the support of the orthogonality measure $d_{q^2}x$ for the big q -Legendre polynomials on $[-1, q^{2\sigma}]$. This can be used to give a different proof for this expression of the Haar functional as a q -integral on $[-1, q^{2\sigma}]$. By determining the spectrum of $\pi(\rho_{\tau,\sigma})$ a similar proof can be given for the Haar functional on the commutative $*$ -algebra generated by $\rho_{\tau,\sigma}$ as an Askey-Wilson integral, see [12] for details. This result is due to Koornwinder [16], who used the corepresentations of $\mathcal{A}_q(SU(2))$ to prove this result.

We use the convention that $v_\lambda(q^\sigma) = 0$ for $\lambda \neq -q^{2n}$, $\lambda \neq q^{2\sigma+2n}$. The eigenvector can be written as $v_\lambda(q^\sigma) = \sum_{n=0}^{\infty} p_n(\lambda) e_n$, where $p_n(\lambda)$ is expressed in terms of the Al-Salam-Carlitz polynomial $U_n^{(a)}$. These polynomials can be considered as the Hermite case

of the big q -Jacobi polynomials. This proposition is proved as follows. The equation $\pi(\rho_{\infty,\sigma}) \sum_{n=0}^{\infty} p_n(\lambda) e_n = \lambda \sum_{n=0}^{\infty} p_n(\lambda) e_n$, $p_{-1}(\lambda) = 0$, $p_0(\lambda) = 1$, leads to a three-term recurrence relation for the required polynomials from which the identification with the Al-Salam–Carlitz polynomials can be made. The corresponding eigenvectors are in $\ell^2(\mathbb{Z}_+)$ only for the given values of λ , which correspond precisely to the discrete mass points in the orthogonality measure for the Al-Salam–Carlitz polynomials. Note that the eigenvectors $v_\lambda(q^\sigma)$ are normalised by the condition $\langle v_\lambda(q^\sigma), e_0 \rangle = p_0(\lambda) = 1$.

The eigenvectors of $\pi(\rho_{\infty,\sigma})$ induce an orthogonal decomposition of the representation space $\ell^2(\mathbb{Z}_+) = V_1^\sigma \oplus V_2^\sigma$, where V_1^σ is the space spanned by the vectors $v_{-q^{2n}}(q^\sigma)$, $n \in \mathbb{Z}_+$, and V_2^σ is the space spanned by the vectors $v_{q^{2\sigma+2n}}(q^\sigma)$, $n \in \mathbb{Z}_+$. The eigenvectors $v_\lambda(q^\sigma)$ of $\pi(\rho_{\infty,\sigma})$ can be considered as a generalisation of the standard basis, which are eigenvectors of $\pi(\rho_{\infty,\infty}) = \lim_{\sigma \rightarrow \infty} \pi(\rho_{\infty,\sigma}) = \pi(q^{-1}\beta\gamma)$, cf. (2.2). It is possible to show that the normalised eigenvector $v_{-q^{2n}}(q^\sigma) / \|v_{-q^{2n}}(q^\sigma)\|$ tends to $i^n e_n$ as $\sigma \rightarrow \infty$ and that $v_{q^{2n+2\sigma}}(q^\sigma) / \|v_{q^{2n+2\sigma}}(q^\sigma)\|$ tends to zero as $\sigma \rightarrow \infty$.

This basis of the representation space $\ell^2(\mathbb{Z}_+)$ is well suited for the calculation of the action of the associated spherical elements (2.12) in this representation. In order to show this we go back to the Hopf $*$ -algebra to find suitable expressions for $\rho_{\tau,\sigma}$.

Proposition 3.3. *For $\sigma, \tau \in \mathbb{R} \cup \{\infty\}$ we have in $\mathcal{A}_q(SU(2))$*

$$\begin{aligned} \beta_{\tau+1,\sigma-1} \gamma_{\tau,\sigma} &= 2q^{\tau+\sigma} \rho_{\tau,\sigma} - q^{2\sigma-1} - q^{2\tau+1}, \\ \gamma_{\tau-1,\sigma+1} \beta_{\tau,\sigma} &= 2q^{\tau+\sigma} \rho_{\tau,\sigma} - q^{2\sigma+1} - q^{2\tau-1}, \\ \alpha_{\tau+1,\sigma+1} \delta_{\tau,\sigma} &= 2q^{\tau+\sigma+1} \rho_{\tau,\sigma} + 1 + q^{2\sigma+2\tau+2}, \\ \delta_{\tau-1,\sigma-1} \alpha_{\tau,\sigma} &= 2q^{\tau+\sigma-1} \rho_{\tau,\sigma} + 1 + q^{2\sigma+2\tau-2}. \end{aligned}$$

Proof. This can be proved straightforwardly from (2.14), (2.8) and the commutation relations (2.1). A more conceptual proof uses [11, prop. 6.5, thm. 5.1], which imply that each product on the right hand side is a polynomial in $\rho_{\tau,\sigma}$. The Clebsch-Gordan series, cf. e.g. [11, §4.7], imply that this polynomial is of degree 1. A simple calculation determines the coefficients in this polynomial. $\square$

Corollary 3.4. *The following relations hold in $\mathcal{A}_q(SU(2))$;*

$$\begin{aligned} \alpha_{\tau,\sigma} \rho_{\tau,\sigma} &= \rho_{\tau-1,\sigma-1} \alpha_{\tau,\sigma}, & \beta_{\tau,\sigma} \rho_{\tau,\sigma} &= \rho_{\tau-1,\sigma+1} \beta_{\tau,\sigma}, \\ \gamma_{\tau,\sigma} \rho_{\tau,\sigma} &= \rho_{\tau+1,\sigma-1} \gamma_{\tau,\sigma}, & \delta_{\tau,\sigma} \rho_{\tau,\sigma} &= \rho_{\tau+1,\sigma+1} \delta_{\tau,\sigma}. \end{aligned}$$

In particular,

$$\begin{aligned} \alpha_{\infty,\sigma} \rho_{\infty,\sigma} &= q^2 \rho_{\infty,\sigma-1} \alpha_{\infty,\sigma}, & \beta_{\infty,\sigma} \rho_{\infty,\sigma} &= \rho_{\infty,\sigma+1} \beta_{\infty,\sigma}, \\ \gamma_{\infty,\sigma} \rho_{\infty,\sigma} &= \rho_{\infty,\sigma-1} \gamma_{\infty,\sigma}, & \delta_{\infty,\sigma} \rho_{\infty,\sigma} &= q^{-2} \rho_{\infty,\sigma+1} \delta_{\infty,\sigma}. \end{aligned}$$

Proof. The special case follows from (2.13). The proof of the first statements are all similar. To prove the first, multiply the last equation of proposition 3.3 by $\alpha_{\tau,\sigma}$ and use

the third equation in the left hand side. Cancelling terms proves the first statement of the corollary. $\square$

Remark 3.5. The elements $\alpha_{\infty,\sigma}$, $\beta_{\infty,\sigma}$, $\gamma_{\infty,\sigma}$ and $\delta_{\infty,\sigma}$ are closely related to the algebra corresponding to a quantum 3-sphere described by Noumi and Mimachi [21]. To see this we calculate the commutation relations among these elements, which can be done straightforwardly or by using [11, props. 6.4, 6.5] as in the proof of proposition 3.3. We get, cf. (2.1),

$$\begin{aligned}\alpha_{\tau-1,\sigma+1}\beta_{\tau,\sigma} &= q\beta_{\tau-1,\sigma-1}\alpha_{\tau,\sigma}, & \alpha_{\tau+1,\sigma-1}\gamma_{\tau,\sigma} &= q\gamma_{\tau-1,\sigma-1}\alpha_{\tau,\sigma}, \\ \gamma_{\tau+1,\sigma+1}\delta_{\tau,\sigma} &= q\delta_{\tau+1,\sigma-1}\gamma_{\tau,\sigma}, & \beta_{\tau+1,\sigma+1}\delta_{\tau,\sigma} &= q\delta_{\tau-1,\sigma+1}\beta_{\tau,\sigma}.\end{aligned}$$

There are no easy analogues of the commutation relations for β and γ and for α and δ , but for $\tau = \infty$ we get from proposition 3.3 and the limit transition (2.13)

$$\alpha_{\infty,\sigma+1}\delta_{\infty,\sigma} - \delta_{\infty,\sigma-1}\alpha_{\infty,\sigma} = q\beta_{\infty,\sigma-1}\gamma_{\infty,\sigma} - q^{-1}\gamma_{\infty,\sigma+1}\beta_{\infty,\sigma}.$$

We consider q^σ as a self-adjoint element of $\mathcal{A}_q(SU(2))$, which commutes with α , β , γ and δ . Now we formally adjoin the unitary operator S to this algebra as in [21, §5], i.e. S commutes with α , β , γ and δ , and $Sq^\sigma = qq^\sigma S$, or S is a shift in σ . Now define $\tilde{x} = q^{\frac{1}{2}}S^{-1}\delta_{\infty,\sigma}$, $\tilde{y} = q^{-\frac{1}{2}}S\alpha_{\infty,\sigma}$, $\tilde{u} = q^{-\frac{1}{2}}S\gamma_{\infty,\sigma}$ and $\tilde{v} = q^{\frac{1}{2}}S^{-1}\beta_{\infty,\sigma}$, then these elements satisfy the relations of the quantum 3-sphere of Noumi and Mimachi [21, §0]. The elements c , d and z defined by Noumi and Mimachi [21, §0, (2.15)] correspond to 1, $q^{2\sigma}$ and $-\rho_{\infty,\sigma}$.

Proposition 3.6. *For $\lambda = -q^{2n}$, $\lambda = q^{2\sigma+2n}$, $n \in \mathbb{Z}_+$, we have*

$$\begin{aligned}\pi(\alpha_{\infty,\sigma})v_\lambda(q^\sigma) &= iq^{\frac{1}{2}-\sigma}(1+\lambda)v_{\lambda/q^2}(q^{\sigma-1}), & \pi(\beta_{\infty,\sigma})v_\lambda(q^\sigma) &= iq^{\frac{1}{2}}(q^{2\sigma}-\lambda)v_\lambda(q^{\sigma+1}), \\ \pi(\gamma_{\infty,\sigma})v_\lambda(q^\sigma) &= iq^{\frac{1}{2}}v_\lambda(q^{\sigma-1}), & \pi(\delta_{\infty,\sigma})v_\lambda(q^\sigma) &= -iq^{\frac{1}{2}+\sigma}v_{\lambda q^2}(q^{\sigma+1}).\end{aligned}$$

Proof. First observe that the result for $\pi(\gamma_{\infty,\sigma})$, respectively $\pi(\delta_{\infty,\sigma})$, implies the result for $\pi(\beta_{\infty,\sigma})$, respectively $\pi(\alpha_{\infty,\sigma})$, by propositions 3.3 and 3.1.

Consider $\pi(\gamma_{\infty,\sigma})$, then we see that $\pi(\gamma_{\infty,\sigma})v_\lambda(q^\sigma) = Cv_\lambda(q^{\sigma-1})$ for some constant C by corollary 3.4 and proposition 3.1. To calculate C we use $\langle v_\lambda(q^\sigma), e_0 \rangle = 1$ and so

$$C = \langle \pi(\gamma_{\infty,\sigma})v_\lambda(q^\sigma), e_0 \rangle = \langle v_\lambda(q^\sigma), -\pi(\beta_{\infty,\sigma-1})e_0 \rangle = iq^{\frac{1}{2}}$$

by (2.15), (2.14) and (2.3).

Similarly, $\pi(\delta_{\infty,\sigma})v_\lambda(q^\sigma) = Cv_{\lambda q^2}(q^{\sigma+1})$ for some constant C with

$$C = \langle \pi(\delta_{\infty,\sigma})v_\lambda(q^\sigma), e_0 \rangle = \langle v_\lambda(q^\sigma), q^{-1}\pi(\alpha_{\infty,\sigma+1})e_0 \rangle = -iq^{\sigma+\frac{1}{2}},$$

which proves the proposition. $\square$

Remark 3.7. If we use $v_\lambda(q^\sigma) = \sum_{n=0}^{\infty} p_n(\lambda)e_n$ and we work out what proposition 3.6 means for the Al-Salam–Carlitz polynomials, which can be expressed as a ${}_2\varphi_1$ -series, then we see that the expressions in proposition 3.6 are equivalent to one of Heine's contiguous relations for the ${}_2\varphi_1$ -series, cf. [6, Ex. 1.9(ii)].

Since $-q\pi(\gamma) = \pi(\beta)$ implies $\pi(\rho_{\tau,\sigma}) = \pi(\rho_{\sigma,\tau})$, $\pi(\alpha_{\tau,\sigma}) = \pi(\alpha_{\sigma,\tau})$, $\pi(\beta_{\tau,\sigma}) = \pi(\gamma_{\sigma,\tau})$, $\pi(\gamma_{\tau,\sigma}) = \pi(\beta_{\sigma,\tau})$ and $\pi(\delta_{\tau,\sigma}) = \pi(\delta_{\sigma,\tau})$, we obtain the following proposition as a corollary of propositions 3.1 and 3.6.

Proposition 3.8. $\ell^2(\mathbb{Z}_+)$ has an orthogonal basis of eigenvectors $v_\lambda(q^\tau)$, where $\lambda = -q^{2n}$, $n \in \mathbb{Z}_+$, $\lambda = q^{2\tau+2n}$, $n \in \mathbb{Z}_+$, for the eigenvalue λ of the self-adjoint operator $\pi(\rho_{\tau,\infty})$. For $\lambda = -q^{2n}$, $\lambda = q^{2\tau+2n}$, $n \in \mathbb{Z}_+$, we have

$$\begin{aligned}\pi(\alpha_{\tau,\infty})v_\lambda(q^\tau) &= iq^{\frac{1}{2}-\tau}(1+\lambda)v_{\lambda/q^2}(q^{\tau-1}), & \pi(\beta_{\tau,\infty})v_\lambda(q^\tau) &= iq^{\frac{1}{2}}v_\lambda(q^{\tau-1}), \\ \pi(\gamma_{\tau,\infty})v_\lambda(q^\tau) &= iq^{\frac{1}{2}}(q^{2\tau}-\lambda)v_\lambda(q^{\tau+1}), & \pi(\delta_{\tau,\infty})v_\lambda(q^\tau) &= -iq^{\frac{1}{2}+\tau}v_{\lambda q^2}(q^{\tau+1}).\end{aligned}$$

It follows from propositions 3.6, 3.8 and (2.12) that for $n \geq 0$

$$(3.1) \quad \begin{aligned}\pi(b_{n,0}^l(\infty, \sigma))v_\lambda(q^\sigma) &= C_n(\sigma)q^{n\sigma}P_{l-n}^{(n,n)}(\lambda; q^{2\sigma}, 1; q^2)v_{\lambda q^{2n}}(q^\sigma), \\ \pi(D.b_{0,n}^l(\tau, \infty))v_\mu(q^\tau) &= C_n(\tau)q^{n(\tau+1)}P_{l-n}^{(n,n)}(\mu; q^{2\tau}, 1; q^2)v_{\mu q^{2n}}(q^\tau)\end{aligned}$$

and

$$(3.2) \quad \begin{aligned}\pi(b_{-n,0}^l(\infty, \sigma))v_\lambda(q^\sigma) &= C_n(\sigma)(-1)^n q^{n(\sigma-1)}(-\lambda, \lambda q^{-2\sigma}; q^{-2})_n \\ &\quad \times P_{l-n}^{(n,n)}(\lambda q^{-2n}; q^{2\sigma}, 1; q^2)v_{\lambda q^{-2n}}(q^\sigma), \\ \pi(D.b_{0,-n}^l(\tau, \infty))v_\mu(q^\tau) &= C_n(\tau)(-1)^n q^{n(\tau-2)}(-\mu, \mu q^{-2\tau}; q^{-2})_n \\ &\quad \times P_{l-n}^{(n,n)}(\mu q^{-2n}; q^{2\tau}, 1; q^2)v_{\mu q^{-2n}}(q^\tau),\end{aligned}$$

where we used $D.\rho_{\tau,\infty} = \rho_{\tau,\infty}$, $D.\alpha_{\tau,\infty} = q^{-\frac{1}{2}}\alpha_{\tau,\infty}$, $D.\beta_{\tau,\infty} = q^{\frac{1}{2}}\beta_{\tau,\infty}$, $D.\gamma_{\tau,\infty} = q^{-\frac{1}{2}}\gamma_{\tau,\infty}$ and $D.\delta_{\tau,\infty} = q^{\frac{1}{2}}\delta_{\tau,\infty}$ and $P_n^{(\alpha,\beta)}(Ax; Ac, AD; q) = P_n^{(\alpha,\beta)}(x; c, D; q)$ for $A > 0$.

From (3.1) and (3.2) it follows that the representation operators corresponding to the right hand side of (2.7) preserve the orthogonal decomposition $\ell^2(\mathbb{Z}_+) \hat{\otimes} \ell^2(\mathbb{Z}_+) = \bigoplus_{i,j=1}^2 V_i^\tau \hat{\otimes} V_j^\sigma$.

Remark 3.9. It is also possible to determine the spectrum of the self-adjoint operator $\pi(\rho_{\tau,\sigma})$ and to use generalised eigenvectors of $\pi(\rho_{\tau,\sigma})$ in the representation space $\ell^2(\mathbb{Z}_+)$. It is then possible to prove an analogue of proposition 3.6 and to calculate the action of the associated spherical elements as in (3.1) and (3.2), but it turns out that only for the choices of the parameters as in this section the orthogonal basis is preserved under the action of the associated spherical elements. This means that restriction $\mu = \infty$ in (2.7) is necessary in order to use the method of this paper to convert (2.7) into an addition formula in commuting variables.

4. ADDITION FORMULA

In this section we derive an explicit addition formula for a two-parameter set of Askey-Wilson polynomials. We show that two known addition formulas can be obtained as limit cases.

From (3.1) and (3.2) we see how the right hand side of (2.7) acts in the basis $v_\mu(q^\tau) \otimes v_\lambda(q^\sigma)$ of the representation space $\ell^2(\mathbb{Z}_+) \hat{\otimes} \ell^2(\mathbb{Z}_+)$. So we now consider the left hand side of (2.7). Since Δ is a $*$ -algebra homomorphism, the left hand side of (2.7) is a polynomial

in $\Delta(\rho_{\tau,\sigma})$. Now we study the self-adjoint operator $(\pi \otimes \pi)\Delta(\rho_{\tau,\sigma})$ in $\ell^2(\mathbb{Z}_+) \hat{\otimes} \ell^2(\mathbb{Z}_+)$. First we consider

$$(4.1) \quad \begin{aligned} \Delta(\beta_{\tau+1,\sigma-1}\gamma_{\tau,\sigma}) &= (D.\alpha_{\tau+1,\infty})(D.\gamma_{\tau,\infty}) \otimes \beta_{\infty,\sigma-1}\alpha_{\infty,\sigma} + \\ &\quad (D.\alpha_{\tau+1,\infty})(D.\delta_{\tau,\infty}) \otimes \beta_{\infty,\sigma-1}\gamma_{\infty,\sigma} + (D.\beta_{\tau+1,\infty})(D.\gamma_{\tau,\infty}) \otimes \delta_{\infty,\sigma-1}\alpha_{\infty,\sigma} \\ &\quad + (D.\beta_{\tau+1,\infty})(D.\delta_{\tau,\infty}) \otimes \delta_{\infty,\sigma-1}\gamma_{\infty,\sigma}, \end{aligned}$$

because we can apply (2.6) for $l = \frac{1}{2}$, $\mu = \infty$ twice, since the elements in (2.14) are generalised matrix elements. We obtain, cf. propositions 3.3, 3.6 and 3.8

$$(4.2) \quad \begin{aligned} 2(\pi \otimes \pi)\Delta(\rho_{\tau,\sigma})v_\mu(q^\tau) \otimes v_\lambda(q^\sigma) &= \\ q^2 v_{\mu q^2}(q^\tau) \otimes v_{\lambda q^2}(q^\sigma) + q^{-2}(1+\lambda)(1+\mu)(1-\lambda q^{-2\sigma})(1-\mu q^{-2\tau})v_{\mu q^{-2}}(q^\tau) \otimes v_{\lambda q^{-2}}(q^\sigma) \\ &\quad + \left(\lambda q^{1-\sigma}(q^{-\tau} - q^\tau) + \mu q^{1-\tau}(q^{-\sigma} - q^\sigma) + \lambda \mu q^{1-\tau-\sigma}(1+q^2) \right) v_\mu(q^\tau) \otimes v_\lambda(q^\sigma). \end{aligned}$$

It follows from (4.2) that $(\pi \otimes \pi)\Delta(\rho_{\tau,\sigma})$ preserves the orthogonal decomposition of the representation space $\ell^2(\mathbb{Z}_+) \hat{\otimes} \ell^2(\mathbb{Z}_+) = \bigoplus_{i,j=1}^2 V_i^\tau \hat{\otimes} V_j^\sigma$. Moreover, it leaves the space $\sum_{m=0}^\infty \mathbb{C} v_{\mu q^{2m}}(q^\tau) \otimes v_{\lambda q^{2m}}(q^\sigma)$ invariant whenever $\mu \in \{-1, q^{2\tau}\}$ or $\lambda \in \{-1, q^{2\sigma}\}$. In each $V_i^\tau \otimes V_j^\sigma$ we have two of such spaces, say $W_{i,j}^\tau$ and $W_{i,j}^\sigma$, where $W_{i,j}^\tau$ is the subspace corresponding to $\mu \in \{-1, q^{2\tau}\}$. In each $W_{i,j}^\nu$, $i, j = 1, 2$, $\nu = \sigma, \tau$, there are $(\pi \otimes \pi)\Delta(\rho_{\tau,\sigma})$ -invariant subspaces. We consider the subspace $W_{1,1}^{\tau,p} = \sum_{m=0}^\infty \mathbb{C} v_{-q^{2m}}(q^\tau) \otimes v_{-q^{2m+2p}}(q^\sigma)$, which is invariant under $(\pi \otimes \pi)\Delta(\rho_{\tau,\sigma})$ for every $p \in \mathbb{Z}_+$.

So we define w_m to be the normalised vector $v_{-q^{2m}}(q^\tau) \otimes v_{-q^{2m+2p}}(q^\sigma)$, i.e. by propositions 3.1 and 3.8,

$$(4.3) \quad w_m = \frac{q^{2m+p} v_{-q^{2m}}(q^\tau) \otimes v_{-q^{2m+2p}}(q^\sigma)}{\sqrt{(q^2, -q^{2-2\tau}; q^2)_m (q^2, -q^{2-2\sigma}; q^2)_{m+p} (-q^{2\tau}, -q^{2\sigma}; q^2)_\infty}}.$$

Then we find the following expression for $(\pi \otimes \pi)\Delta(\rho_{\tau,\sigma})$;

$$(4.4) \quad \begin{aligned} 2(\pi \otimes \pi)\Delta(\rho_{\tau,\sigma})w_m &= a_m w_{m+1} + b_m w_m + a_{m-1} w_{m-1}, \\ a_m &= \sqrt{(1 - q^{2m+2})(1 - q^{2m+2p+2})(1 + q^{2-2\tau+2m})(1 + q^{2+2p-2\sigma+2m})}, \\ b_m &= q^{2m} \left(q^{1+2p-\sigma}(q^\tau - q^{-\tau}) + q^{1-\tau}(q^\sigma - q^{-\sigma}) + q^{2m+2p+1-\tau-\sigma}(1+q^2) \right). \end{aligned}$$

The three-term recurrence for orthonormal polynomials corresponding to (4.4) can be solved in terms of Askey-Wilson polynomials, or in terms of the q -Laguerre case of the q -Jacobi polynomials defined in (2.10). So we define

$$(4.5) \quad l_n^{(\alpha)}(x; s, t \mid q) = p_n(x; q^{1/2+\alpha}s/t, q^{1/2}t/s, -q^{1/2}/(st), 0 \mid q),$$

i.e. by letting $\beta \rightarrow \infty$ in (2.10). From the three-term recurrence relation for the Askey-Wilson polynomials, cf. Askey and Wilson [3, §1], we obtain

$$\begin{aligned} 2xl_n(x) &= l_{n+1}(x) + (1 - q^n)(1 - q^{\alpha+n})(1 + q^n s^{-2})(1 + q^{n+\alpha} t^{-2})l_{n-1}(x) \\ &\quad + q^n \left((t - t^{-1})q^{\frac{1}{2}} s^{-1} + (s - s^{-1})q^{\frac{1}{2}+\alpha} t^{-1} + (1 + q)q^{\frac{1}{2}+n+\alpha} s^{-1} t^{-1} \right) l_n(x), \end{aligned}$$

where $l_n(x) = l_n^{(\alpha)}(x; s, t \mid q)$. The orthogonality measure for the q -Laguerre polynomials follows from [3, thm. 2.4], and as in [16, (2.7)] we denote the normalised orthogonality measure by $dm^{(\alpha)}(\cdot; s, t \mid q)$.

If we denote the corresponding orthonormal polynomials by $\tilde{l}_n^{(\alpha)}(x; s, t \mid q)$, we see that the three-term recurrence (4.4) is solved by the q -Laguerre polynomials $\tilde{l}_m^{(p)}(\cdot; q^\tau, q^\sigma \mid q^2)$. The spectral theory of Jacobi matrices shows that the spectral decomposition E of the self-adjoint operator $(\pi \otimes \pi)\Delta(\rho_{\tau, \sigma})$ on the Hilbert space $W_{1,1}^{\tau, p}$ with orthonormal basis w_m , $m \in \mathbb{Z}_+$, is given by

$$\langle E(B)w_n, w_m \rangle = \int_B (\tilde{l}_n^{(p)} \tilde{l}_m^{(p)})(x; q^\tau, q^\sigma \mid q^2) dm^{(p)}(x; q^\tau, q^\sigma \mid q^2)$$

for B a Borel subset of $\mathbb{R}$, cf. Berezanskii [4, Ch. VII, §1], Dombrowski [5]. The mapping $\Lambda: W_{1,1}^{\tau, p} \rightarrow L^2(\mathbb{R}, dm^{(p)}(x; q^\tau, q^\sigma \mid q^2))$ given by $\Lambda(w_m)(x) = \tilde{l}_m^{(p)}(x; q^\tau, q^\sigma \mid q^2)$ extends to a unitary mapping, since the corresponding moment problem is determined. Λ intertwines $(\pi \otimes \pi)\Delta(\rho_{\tau, \sigma})$ on $W_{1,1}^{\tau, p}$ with the multiplication operator M , $Mf(x) = xf(x)$, on this L^2 -space.

Next we apply $\Lambda \circ p_l^{(0,0)}((\pi \otimes \pi)\Delta(\rho_{\tau, \sigma}); q^\tau, q^\sigma \mid q^2)$ to w_m given by (4.3) on the one hand. On the other hand we apply $\pi \otimes \pi$ of the right hand side of (2.7) for $\mu = \infty$ to the vector w_m for which we use (3.1) and (3.2) before we apply Λ . This results in the addition formula for the two-parameter q -Legendre polynomial $p_l^{(0,0)}(x; q^\sigma, q^\tau \mid q^2)$ as an identity in $L^2(\mathbb{R}, dm^{(p)}(x; q^\tau, q^\sigma \mid q^2))$, but since it only involves polynomials it holds everywhere. The following theorem is then obtained after replacing q^2 by q and τ, σ by $2\tau, 2\sigma$.

Theorem 4.1. *The following addition formula holds for $l, m, p \in \mathbb{Z}_+$, $\sigma, \tau \in \mathbb{R}$, $x \in \mathbb{C}$:*

$$\begin{aligned} p_l^{(0,0)}(x; q^\tau, q^\sigma \mid q) l_m^{(p)}(x; q^\tau, q^\sigma \mid q) = \\ \sum_{n=0}^l D^{n,l}(\tau, \sigma) P_{l-n}^{(n,n)}(-q^m; q^{2\tau}, 1; q) P_{l-n}^{(n,n)}(-q^{m+p}; q^{2\sigma}, 1; q) l_{m+n}^{(p)}(x; q^\tau, q^\sigma \mid q) \\ + \sum_{n=1}^l D^{n,l}(\tau, \sigma) (q^m, q^{m+p}, -q^{m+p-2\sigma}, -q^{m-2\tau}; q^{-1})_n \\ \times P_{l-n}^{(n,n)}(-q^{m-n}; q^{2\tau}, 1; q) P_{l-n}^{(n,n)}(-q^{m+p-n}; q^{2\sigma}, 1; q^2) l_{m-n}^{(p)}(x; q^\tau, q^\sigma \mid q), \end{aligned}$$

with the constant given by

$$D^{n,l}(\tau, \sigma) = (-q^{2\sigma-l}, -q^{2\tau-l}; q)_{l-n} \frac{(q^{l-n+1}; q)_n}{(q; q)_n} (q^{n+1}; q)_l q^{\frac{1}{2}(l-n)(l-n-2\sigma-2\tau)}$$

and the notation for the q -Legendre polynomials $p_l^{(0,0)}(\cdot; q^\tau, q^\sigma \mid q)$, the q -Laguerre polynomials $l_m^{(p)}(\cdot; q^\tau, q^\sigma \mid q)$ and the big q -ultraspherical polynomials $P_{l-n}^{(n,n)}(\cdot; q^{2\tau}, 1; q)$ defined by (2.10), (4.5) and (2.16).

Proof. It remains to calculate the constants involved. If h_m denotes the square norm of $l_m^{(p)}(\cdot; q^\tau, q^\sigma \mid q^2)$, then we use, cf. Askey and Wilson [3, thm. 2.3],

$$\frac{\|v_{-q^{2m+2n}}(q^\tau) \otimes v_{-q^{2m+2n+2p}}(q^\sigma)\|}{\|v_{-q^{2m}}(q^\tau) \otimes v_{-q^{2m+2p}}(q^\sigma)\|} \sqrt{\frac{h_m}{h_{m+n}}} = q^{-2n}$$

to see that before changing q^2 to q and τ, σ to $2\tau, 2\sigma$ we have

$$D^{n,l}(\tau, \sigma) = q^l(q^{2l+2}; q^2)_l C_n(\sigma) C_n(\tau) q^{n(\sigma+\tau-1)}.$$

A straightforward calculation yields the result. $\square$

Remark 4.2. In deriving theorem 4.1 we have chosen a certain generalised eigenvector of $(\pi \otimes \pi)\Delta(\rho_{\tau,\sigma})$ from the space $W_{1,1}^{\tau,p} \subset \ell^2(\mathbb{Z}_+) \hat{\otimes} \ell^2(\mathbb{Z}_+)$. It is also possible to chose a generalised eigenvector from similarly defined spaces $W_{i,j}^{\nu,p}$, $i, j = 1, 2$, $\nu = \sigma, \tau$, so that we obtain in total eight of such eigenvectors. This results in eight of these addition formulas, which are obviously four by four equivalent by interchanging τ and σ and noting that the q -Legendre polynomial involved is also invariant under such a change. The remaining four types of addition formulas differ, since the big q -ultraspherical polynomials are evaluated at other points of the spectrum. However, from

$$(4.6) \quad P_n^{(\alpha,\beta)}(-x; c, d; q) = (-q^{\alpha-\beta}d/c)^n \frac{(q^{\beta+1}, -q^{\beta+1}c/d; q)_n}{(q^{\alpha+1}, -q^{\alpha+1}d/c; q)_n} P_n^{(\beta,\alpha)}(x; d, c; q),$$

which is a direct consequence of the orthogonality relations, cf. [1], and the trivial relation $P_n^{(\alpha,\beta)}(Ax; Ac, Ad; q) = P_n^{(\alpha,\beta)}(x; c, d; q)$, for $A > 0$, we see that

$$(4.7) \quad P_{l-n}^{(n,n)}(-q^m; q^{2\tau}, 1; q) = (-q^{-2\tau})^{l-n} \frac{(-q^{n+1+2\tau}; q)_{l-n}}{(-q^{n+1-2\tau}; q)_{l-n}} P_{l-n}^{(n,n)}(q^{m-2\tau}; q^{-2\tau}, 1; q).$$

If we use this in theorem 4.1 and we next change x into $-x$ using $p_n^{(\alpha,\beta)}(-x; s, t \mid q) = (-1)^n p_n^{(\alpha,\beta)}(x; -s, t \mid q)$ for the q -Legendre polynomial, $\alpha = \beta = 0$, and for the q -Laguerre polynomial, $\beta \rightarrow \infty$, and we change q^τ to $-q^{-\tau}$, then we obtain the same addition formula as if we had started off with the space $W_{2,1}^{\tau,p}$, i.e. with vectors of the type $v_{q^{2\tau+2m}}(q^\tau) \otimes v_{-q^{2m+2p}}(q^\sigma)$, $m \in \mathbb{Z}_+$. A similar approach can be used for the other big q -ultraspherical polynomial, and a combination of both shows that theorem 4.1 contains all the other possibilities by symmetry considerations.

We also obtain the same addition formula if we would have started with one of the other irreducible $*$ -representations of the Hopf $*$ -algebra $\mathcal{A}_q(SU(2))$.

Remark 4.3. Theorem 4.1 contains as special cases two addition formulas for q -Legendre polynomials previously obtained from the quantum $SU(2)$ group. Firstly, we rescale $x = q^{1-\sigma-\tau}y/2$ and we take the limits $\tau \rightarrow \infty$. The limit transition of Askey-Wilson polynomials to big q -Jacobi polynomials as in [16, (6.2)] shows that $p_l^{(0,0)}(x; q^\tau, q^\sigma \mid q)$ and $l_m^{(p)}(x; q^\tau, q^\sigma \mid q)$ tend to the big q -Legendre and big q -Laguerre polynomials. Moreover,

the big q -ultraspherical polynomial $P_{l-n}^{(n,n)}(x; q^{2\tau}, 1; q)$ tends to a little q -Jacobi polynomial, and we obtain an addition formula for a big q -Legendre polynomial. If we next let the order p of the big q -Laguerre polynomial tend to infinity, so that the big q -Laguerre polynomials tend to the Al-Salam–Carlitz $U_n^{(a)}$, we obtain the addition formula for the big q -Legendre polynomials derived in [10, thm. 4.3]. For this we have to observe that a big q -ultraspherical polynomial of argument zero can be rewritten as a dual q -Krawtchouk polynomial by [6, (III.5), (III.6)].

Secondly, rescaling $x = q^{1-\sigma-\tau}y/2$ and letting both $\sigma, \tau \rightarrow \infty$, and using the limit transition of the Askey-Wilson polynomials to the little q -Jacobi polynomial, cf. [16, (6.4)], and the limit transition of the big q -Jacobi polynomials to the little q -Jacobi polynomials, shows that the addition formula in theorem 4.1 contains Koornwinder’s addition formula for the little q -Legendre polynomials [15, thm. 4.1] as a special case, see Rahman [22] for an analytic proof.

Remark 4.4. Another special case of theorem 4.1 is the case $\sigma = \tau = 0$, which gives an addition formula for the continuous q -Legendre polynomial. This addition formula is an alternative for the addition formula for the continuous q -Legendre polynomials derived by Rahman and Verma [23, (1.24) with $a = q^{1/4}$], see also [9, §4].

5. PRODUCT FORMULA

Using the orthogonality relations for the q -Laguerre polynomial a product formula can be easily obtained from the addition formula. For this we need the normalised orthogonality measure $dm^{(\alpha)}(x; s, t \mid q)$ for the q -Laguerre polynomials;

$$(5.1) \quad \int_{\mathbb{R}} \left(l_m^{(\alpha)} l_n^{(\alpha)} \right) (x; s, t \mid q) dm^{(\alpha)}(x; s, t \mid q) = \delta_{n,m}(q, q^{1+\alpha}, -q^{-2}, -q^{1+\alpha}t^{-2} : q)_n,$$

cf. Askey and Wilson [3, thm. 2.5] in combination with (4.5) for the explicit form of the orthogonality measure as an absolutely continuous measure on $[-1, 1]$ plus a finite number, possibly zero, of discrete mass points off $[-1, 1]$.

Now multiply theorem 4.1 by $l_{m+r}^{(p)}(x; q^\tau, q^\sigma \mid q)$ and integrate against the normalised orthogonality measure $dm^{(p)}(x; q^\tau, q^\sigma \mid q^2)$ use the orthogonality relations (5.1) for the q -Laguerre polynomials to obtain the following product formula for the product of two big q -ultraspherical polynomials, which for $n = 0$ gives a q -analogue of the product formula for the Legendre polynomials (1.2).

Corollary 5.1. *For $l, n \in \mathbb{Z}_+$, $0 \leq n \leq l$ we have the product formula*

$$P_{l-n}^{(n,n)}(-q^m; q^{2\tau}, 1; q) P_{l-n}^{(n,n)}(-q^{m+p}; q^{2\sigma}, 1; q) = \frac{1}{C} \int_{\mathbb{R}} p_l^{(0,0)}(x; q^\tau, q^\sigma \mid q) l_m^{(p)}(x; q^\tau, q^\sigma \mid q) l_{m+n}^{(p)}(x; q^\tau, q^\sigma \mid q) dm^{(p)}(x; q^\tau, q^\sigma \mid q),$$

with

$$C = D^{n,l}(\tau, \sigma)(q, q^{1+p}, -q^{1-2\tau}, -q^{1+p-2\sigma}; q)_{m+n}.$$

The same product formula is obtained if we multiply theorem by $l_{m-r}^{(p)}(x; q^\tau, q^\sigma \mid q)$ before using the orthogonality relations. A product formula for big q -ultraspherical polynomials at other points of the spectrum follows from (4.7).

6. THE LIMIT CASE $q \uparrow 1$

We consider the limit transition $q \uparrow 1$ of the addition and product formulas from theorem 4.1 and corollary 5.1 to the addition and product formulas for the Legendre polynomials as presented in §1. This can be done using the theorems of Van Assche and Koornwinder [25, thm. 1, 2], cf. [10, §5] for another example of the use of these theorems.

We divide both sides of the addition formula in theorem 4.1 by $l_m^{(p)}(x; q^\tau, q^\sigma; | q)(q; q)_l$ and we substitute $q = c^{1/m}$, $0 < c < 1$, so that $q \uparrow 1$ corresponds to $m \rightarrow \infty$, and $p = mr$ with $r = \ln p / \ln c \in \mathbb{Z}_+$. Also, put $t = c^r$. Under these identifications we see that the various q -Jacobi polynomials tend to the Jacobi polynomials as $m \rightarrow \infty$, e.g.

$$\begin{aligned} P_{l-n}^{(n,n)}(-q^{m+p-n}; q^{2\sigma}, 1; q) &\rightarrow R_{l-n}^{(n,n)}(-ct), \\ (q; q)_l^{-1} p_l^{(0,0)}(x; q^\tau, q^\sigma | q) &\rightarrow 4^l R_l^{(0,0)}(x). \end{aligned}$$

Also

$$\begin{aligned} \frac{D^{n,l}(\tau, \sigma)}{(q; q)_l} &\rightarrow 4^{l-n} \frac{(l-n+1)_n (n+1)_l}{n! l!}, \\ (q^m, q^{m+p}, -q^{m+p-2\sigma}, -q^{m-2\tau}; q^{-1})_n &\rightarrow ((1-c^2)(1-c^2 t^2))^n. \end{aligned}$$

The non-trivial limit transition of the quotient of two q -Laguerre polynomials is handled using theorem 1 of Van Assche and Koornwinder [25] and we obtain for $k \in \mathbb{Z}$

$$\lim_{m \rightarrow \infty} \frac{l_{m+k}^{(mr)}(x; c^{\tau/m}, c^{\sigma/m} | c^{1/m})}{l_m^{(mr)}(x; c^{\tau/m}, c^{\sigma/m} | c^{1/m})} = ((1-c^2)(1-t^2 c^2))^{k/2} \rho^k((x-c^2 t)/\sqrt{(1-c^2)(1-t^2 c^2)}),$$

where $\rho(x) = x + \sqrt{x^2 - 1}$ ($|\rho(x)| > 1$ for $x \in \mathbb{C} \setminus [-1, 1]$). Using these limit transitions in theorem 4.1 and $2T_n(x) = \rho^k(x) + \rho^{-k}(x)$ shows that for $q \uparrow 1$ we obtain the addition formula (1.1) for the Legendre polynomials.

The limit transition of the product formula of corollary 5.1 to the product formula (1.2) is treated similarly, but now we have to use theorem 2 of Van Assche and Koornwinder [25] on the weak asymptotics of the q -Laguerre polynomials. A straightforward calculation shows that this theorem implies for continuous f and $n \in \mathbb{Z}_+$

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathbb{R}} f(x) \left(\tilde{l}_m^{(mr)} \tilde{l}_{m+n}^{(mr)} \right) (x; q^\tau, q^\sigma | q) dm^{(mr)}(x; q^\tau, q^\sigma | q) = \\ \frac{1}{\pi} \int_{B-2A}^{B+2A} f(x) \frac{T_n((x-B)/(2A))}{\sqrt{4A^2 - (x-B)^2}} dx \end{aligned}$$

with $B = c^2 t$, $2A = \sqrt{(1-c^2)(1-c^2 t^2)}$. Using the previous limit transitions of this section we see that the product formula of corollary 5.1 tends to the classical product formula (1.2) as $q \uparrow 1$, see [25, §4], [10, §5] for similar limit transitions.

REFERENCES

1. G.E. Andrews and R. Askey, *Classical orthogonal polynomials*, “Polynômes Orthogonaux et Applications” (C. Brezinski, A. Draux, A.P. Magnus, P. Maroni, and A. Ronveaux, eds.), Lecture Notes Math. 1171, Springer-Verlag, New York, 1985, pp. 36–62.
2. R. Askey, *Orthogonal Polynomials and Special Functions*, CBMS-NSF Regional Conference Series Applied Math. **21**, SIAM, Philadelphia PA, 1975.
3. R. Askey and J. Wilson, *Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials*, *Memoirs Amer. Math. Soc.* **54** (1985), no. 319.
4. J.M. Berezanskiĭ, *Expansions in Eigenfunctions of Selfadjoint Operators*, Transl. Math. Monographs 17, Amer. Math. Soc., Providenc RI, 1968.
5. J. Dombrowski, *Orthogonal polynomials and functional analysis*, “Orthogonal Polynomials: Theory and Practice” (P. Nevai, ed.), NATO ASI series C, vol. 294, Kluwer, Dordrecht, 1990, pp. 147–161.
6. G. Gasper and M. Rahman, *Basic Hypergeometric Series*, Encyclopedia of Mathematics and its Applications 35, Cambridge University Press, 1990.
7. M. Jimbo, *A q -difference analogue of $U(g)$ and the Yang-Baxter equation*, *Lett. Math. Phys.* **10** (1985), 63–69.
8. E.G. Kalnins and W. Miller, *Models of q -algebra representations: q -integral transforms and addition theorems*, *J. Math. Phys.* **35** (1994), 1951–1975.
9. H.T. Koelink, *The addition formula for continuous q -Legendre polynomials and associated spherical elements on the $SU(2)$ quantum group related to Askey-Wilson polynomials*, *SIAM J. Math. Anal.* **25** (1994), 197–217.
10. ———, *Addition formula for big q -Legendre polynomials from the quantum $SU(2)$ group*, *Canad. J. Math.* (to appear).
11. ———, *Askey-Wilson polynomials and the quantum $SU(2)$ group: survey and applications*, *Acta Appl. Math.* (to appear).
12. H.T. Koelink and J. Verding, *Spectral analysis and the Haar functional on the quantum $SU(2)$ group*, in preparation.
13. T.H. Koornwinder, *Representations of the twisted $SU(2)$ quantum group and some q -hypergeometric orthogonal polynomials*, *Proc. Kon. Ned. Akad. van Wetensch., Ser. A* **92** (Indag. Math. **51**) (1989), 97–117.
14. ———, *Orthogonal polynomials in connection with quantum groups*, “Orthogonal Polynomials: Theory and Practice” (P. Nevai, ed.), NATO ASI series C, vol. 294, Kluwer, Dordrecht, 1990, pp. 257–292.
15. ———, *The addition formula for little q -Legendre polynomials and the $SU(2)$ quantum group*, *SIAM J. Math. Anal.* **22** (1991), 295–301.
16. ———, *Askey-Wilson polynomials as zonal spherical functions on the $SU(2)$ quantum group*, *SIAM J. Math. Anal.* **24** (1993), 795–813.
17. T. Masuda, K. Mimachi, Y. Nakagami, M. Noumi and K. Ueno, *Representations of the quantum group $SU_q(2)$ and the little q -Jacobi polynomials*, *J. Funct. Anal.* **99** (1991), 357–386.
18. M. Noumi, *Quantum groups and q -orthogonal polynomials. Towards a realization of Askey-Wilson polynomials on $SU_q(2)$* , “Special Functions” (M. Kashiwara and T. Miwa, eds.), ICM-90 Satellite Conference Proceedings, Springer-Verlag, New York, 1991, pp. 260–288.
19. M. Noumi and K. Mimachi, *Askey-Wilson polynomials and the quantum group $SU_q(2)$* , *Proc. Japan Acad., Ser. A* **66** (1990), 146–149.
20. ———, *Askey-Wilson polynomials as spherical functions on $SU_q(2)$* , “Quantum Groups” (P.P. Kulish, ed.), Lecture Notes Math. 1510, Springer-Verlag, New York, 1992, pp. 98–103.
21. ———, *Spherical functions on a family of quantum 3-spheres*, *Compositio Math.* **83** (1992), 19–42.
22. M. Rahman, *A simple proof of Koornwinder’s addition formula for the little q -Legendre polynomials*, *Proc. Amer. Math. Soc.* **107** (1989), 373–381.
23. M. Rahman and A. Verma, *Product and addition formulas for the continuous q -ultraspherical polynomials*, *SIAM J. Math. Anal.* **17** (1986), 1461–1474.
24. L.L. Vaksman and Y.S. Soibelman, *Algebra of functions on the quantum group $SU(2)$* , *Funct. Anal. Appl.* **22** (1988), 170–181.

- 25. W. Van Assche and T.H. Koornwinder, *Asymptotic behaviour for Wall polynomials and the addition formula for little q -Legendre polynomials*, SIAM J. Math. Anal. **22** (1991), 302–311.
- 26. S.L. Woronowicz, *Twisted $SU(2)$ group. An example of non-commutative differential calculus*, Publ. Res. Inst. Math. Sci., Kyoto Univ. **23** (1987), 117–181.
- 27. ———, *Compact matrix pseudo groups*, Commun. Math. Phys. **111** (1987), 613–665.

DEPARTEMENT WISKUNDE, KATHOLIEKE UNIVERSITEIT LEUVEN, CELESTIJNENLAAN 200 B, B-3001
LEUVEN (HEVERLEE), BELGIUM

E-mail address: `erik.koelink@wis.kuleuven.ac.be`